

Supplemental Material 2 - Excluded articles and reasons for exclusion (n= 16)

Author, Year	Reasons for exclusion
Ahmadvazdeh et al., 2017	3
Artieres, 1978	1
Costa Branco Neves, 2005	6
Da Silva et al., 2015	3
Favrelière et al., 2020	4
Gaber et al., 2014	3
Galarza-Delgado et al., 2018	3
Gamal et al., 2021	3
Gündüz et al., 2019	3
Kastanioudakis et al., 1995	3
Levy et al., 1991	4
McGready et al., 2002	4
Nott et al., 2020	2
Patatt et al., 2021	3
Silva et al., 2015	2
Tumwikirize et al., 2011	5

1- Study not found (n=1); 2- Congress abstract (n=2); 3- Analysis does not separate exposed and unexposed groups (n=8); 4- Does not mention which tests performed for audiological evaluation (n=3); 5- It does not mention how many participants who had hearing loss took the medication (n=1); 6- Includes participants with diseases associated with hearing loss (n=1)

REFERENCES

- Ahmadvazdeh, A., Daraei, M., Jalessi, M., Peyvandi, A. A., Amini, E., Ranjbar, L. A., & Daneshi, A. (2017). Hearing status in patients with rheumatoid arthritis. *Journal of Laryngology and Otology*, 131(10), 895-899. <https://doi.org/10.1017/S0022215117001670>
- Artieres, J. (1978). Therapeutic effects of Tanakan (chloroquine) on hearing loss and symptoms of auditory disturbance. *Lyon Mediterraneo Medical*, 14(15), 2503-2515. <https://www.scopus.com/inward/record.uri?eid=2-s2.0-0018081880&partnerID=40&md5=e8867f89741d511ca7117891b978400a>
- Costa Branco Neves, M. V. (2005). *Estudio de los efectos ototóxicos en 725 pacientes tratados con antimaláricos en el hospital central de Maputo (Mozambique)*. Universitat Autònoma de Barcelona.
- da Silva, V. B., Sousa, M. A., Kuniyoshi, I. C., Aurélio, F. S., Sampaio, A. L. L., & De Oliveira, C. A. C. P. (2015). Risk of hearing alterations in newborns of mothers treated for malaria. *International Tinnitus Journal*, 19(2), 20-25. <https://doi.org/10.5935/0946-5448.20150004>
- Favrelière, S., Delaunay, P., Lebreton, J. P., Rouby, F., Atzenhoffer, M., Lafay-Chebassier, C., & Péault-Pochat, M. C. (2020). Drug-induced hearing loss: a case/non-case study in the French pharmacovigilance database. *Fundamental and Clinical Pharmacology*. <https://doi.org/10.1111/fcp.12533>
- Gaber, W., Ezzat, Y., El Fayoumy, N. M., Helmy, H., & Mohey, A. M. (2014). Detection of asymptomatic cranial neuropathies in patients with systemic lupus erythematosus and their relation to antiribosomal P antibody levels and disease activity. *Clin Rheumatol*, 33(10), 1459-1466. <https://doi.org/10.1007/s10067-014-2679-y>
- Galarza-Delgado, D. A., Gonzalez, M. J. V., Torres, J. R., Soto-Galindo, G. A., Flores, L. M., & Gonzalez, J. L. T. (2018). Early hearing loss detection in rheumatoid arthritis and

- primary Sjogren syndrome using extended high frequency audiometry. *Clin Rheumatol*, 37(2), 367-373. <https://doi.org/10.1007/s10067-017-3959-0>
- Gamal, N. M., El-Oseily, A. M. A., Mostafa, H., Badawy, E. R., & Abd Elsamea, M. H. (2021). Detection of hearing loss in rheumatoid arthritis patients using extended high frequency audiometry: Is it related to disease activity and severity? *The Egyptian Rheumatologist*, 43(3), 219-224.
- Gündüz, B., Yıldırım, N., Güven, S. C., Orhan, E., Karamert, R., & Günendi, Z. (2019). Evaluation of medial olivocochlear efferent system and hearing loss in patients with primary Sjögren's syndrome. *Turkish Journal of Medical Sciences*, 49(6), 1647-1654. <https://doi.org/10.3906/sag-1901-128>
- Kastanioudakis, I., Skevas, A., Danielidis, V., Tsiaikou, E., Drosos, A. A., & Moustopoulos, M. H. (1995). Inner ear involvement in rheumatoid arthritis: a prospective clinical study. *J Laryngol Otol*, 109(8), 713-718. <https://doi.org/10.1017/s0022215100131135>
- Levy, M., Buskila, D., Gladman, D. D., Urowitz, M. B., & Koren, G. (1991). Pregnancy outcome following first trimester exposure to chloroquine. *Am J Perinatol*, 8(3), 174-178. <https://doi.org/10.1055/s-2007-999371>
- McGready, R., Thwai, K. L., Cho, T., Samuel, Looareesuwan, S., White, N. J., & Nosten, F. (2002). The effects of quinine and chloroquine antimalarial treatments in the first trimester of pregnancy. *Trans R Soc Trop Med Hyg*, 96(2), 180-184. [https://doi.org/10.1016/S0035-9203\(02\)90297-X](https://doi.org/10.1016/S0035-9203(02)90297-X)
- Nott, K., Nott, V., & Compeyrot-Lacassagne, S. (2020). *Paediatric Sarcoidosis: Phenotype of a Retrospective Cohort of Biopsy-proven Patients* ACR Convergence 2020, Virtual. <https://acrabstracts.org/abstract/paediatric-sarcoidosis-phenotype-of-a-retrospective-cohort-of-biopsy-proven-patients/>
- Patatt, F. S. A., Sampaio, A. L. L., Taui, P. L., & Oliveira, C. A. C. P. d. (2021). Hearing of neonates without risk indicators for hearing loss and use of antimalarial drugs during pregnancy: a historical cohort study in the Northern Region of Brazil. *Braz J Otorhinolaryngol*, 87, 19-27.
- Silva, V. B., Sousa, M. A., Kuniyoshi, I. C., Aurélio, F. S., Sampaio, A. L., & Oliveira, C. A. (2015). Risk of hearing alterations in newborns of mothers treated for malaria. *Int Tinnitus J*, 19(2), 20-25. <https://doi.org/10.5935/0946-5448.20150004>
- Tumwikirize, W. A., Ogwale-Okeng, J. W., Vernby, A., Anokbonggo, W. W., Gustafsson, L. L., & Lundborg, S. C. (2011). Adverse drug reactions in patients admitted on Internal Medicine wards in a district and Regional Hospital in Uganda. *African Health Sciences*, 11(1), 72-78. <https://www.scopus.com/inward/record.uri?eid=2-s2.0-79953804322&partnerID=40&md5=8740e9da277f1e3e047059b9d812109b>